

Package: FMsmnReg (via r-universe)

November 5, 2024

Type Package

Title Regression Models with Finite Mixtures of Skew Heavy-Tailed Errors

Version 1.0

Date 2016-03-29

Author Luis Benites Sanchez and Rocio Paola Maehara and Victor Hugo Lachos

Maintainer Luis Benites Sanchez <lbenitesanchez@gmail.com>

Imports mvtnorm

Description Fit linear regression models where the random errors follow a finite mixture of of Skew Heavy-Tailed Errors.

License GPL (>= 2)

NeedsCompilation no

Date/Publication 2016-03-30 08:14:20

Repository <https://lbenitesanchez.r-universe.dev>

RemoteUrl <https://github.com/lbenitesanchez/fmsmsnreg>

RemoteRef HEAD

RemoteSha ad304f7524fa85d88c3842867afbc068dfdbbeff

Contents

FMsmnReg-package	2
ais	3
FMsmnReg	4
horses	6
Index	8

FMsmnsReg-package *Linear Regression Models with Finite Mixtures of Skew Heavy-Tailed Errors*

Description

This package contains a principal function that performs to estimate the parameters of a regression model considering an error that follows a finite mixture of Skew Heavy-Tailed Errors, using an analytically simple and efficient EM-type algorithm for iteratively computing maximum likelihood estimates of the parameters.

Details

Package: FMsmnsReg
Type: Package
Version: 1.0
Date: 2016-03-30
License: GPL (>=2)

Author(s)

Luis Benites Sanchez <lbenitesanchez@gmail.com>, Rocio Paola Maehara <rmaeharaa@gmail.com>
and Victor Hugo Lachos <hlachos@ime.unicamp.br>

References

Basso, R. M., Lachos, V. H., Cabral, C. R., Ghosh, P., 2010. Robust mixture modeling based on scale mixtures of skew - normal distributions. *Computational Statistics & Data Analysis*.

Lachos, V. H., Ghosh, P., Arellano-Valle, R. B., 2010. Likelihood based inference for skew-normal independent linear mixed models. *Statistica Sinica* 20, 303 - 322.

See Also

[FMsmnsReg](#)

Examples

#See examples for the FMsmnsReg function linked above.

 ais

Australian institute of sport data

Description

Data on 102 male and 100 female athletes collected at the Australian Institute of Sport.

Format

This data frame contains the following columns:

Sex (0 = male or 1 = female)

Ht height (cm)

Wt weight (kg)

LBM lean body mass

RCC red cell count

WCC white cell count

Hc Hematocrit

Hg Hemoglobin

Ferr plasma ferritin concentration

BMI body mass index, weight/height²

SSF sum of skin folds

Bfat Percent body fat

Label Case Labels

Sport Sport

References

S. Weisberg (2005). *Applied Linear Regression*, 3rd edition. New York: Wiley, Section 6.4

Examples

```
##Load the data
library(FMsmnReg)
data(ais)
attach(ais)
#Set the response y and covariate x
x1  <- cbind(1,SSF,Ht)
y   <- Bfat

##Fits a linear Regression Model with Finite Mixtures of Skew t
parST <- FMsmnReg(y, x1, g=2, get.init = TRUE, criteria = TRUE,
  group = FALSE,family = "Skew.t", error = 10^-4,
  iter.max = 2000,obs.prob= FALSE, kmeans.param = NULL,show.converge=FALSE,cp=0.5)
```

```
##Fits a linear Regression Models with Finite Mixtures of Skew normal
parSN <- FMsmnReg(y, x1, g=2, get.init = TRUE, criteria = TRUE,
  group = FALSE,family = "Skew.normal", error = 10^-4,
  iter.max = 5000,obs.prob= FALSE, kmeans.param = NULL,show.converge=FALSE,cp=0.5)

##Fits a linear Regression Models with Finite Mixtures of Skew Contaminated Normal
parCN <- FMsmnReg(y, x1, g=2, get.init = TRUE, criteria = TRUE,
  group = FALSE,family = "Skew.cn", error = 10^-4,
  iter.max = 5000,obs.prob= FALSE, kmeans.param = NULL,show.converge=FALSE,cp=0.5)
```

FMsmnReg	<i>Linear Regression Models with Finite Mixtures of Skew Heavy-Tailed Errors</i>
----------	--

Description

Performs a Finite Mixture of Scale Mixture Skew Normal Regression Model using EM-type algorithm (ECME) for iteratively computing maximum likelihood estimates of the parameters.

Usage

```
FMsmnReg(y, x1, Abetas = NULL, medj= NULL, sigma2 = NULL, shape = NULL,
  pii = NULL, g = NULL, get.init = TRUE, criteria = TRUE, group = FALSE,
  family = "Skew.normal", error = 0.00001, iter.max = 100, obs.prob= FALSE,
  kmeans.param = NULL, show.convergence=TRUE, cp=0.4)
```

Arguments

y	the response matrix (dimension nx1)
x1	Matrix or vector of covariates.
Abetas	Parameters of vector regression dimension ($p + 1$) include intercept
medj	a list of g arguments of vectors of initial values (dimension p) for the location parameters
sigma2	a list of g arguments of matrices of initial values (dimension pxp) for the scale parameters
shape	a list of g arguments of vectors of initial values (dimension p) for the skewness parameters
pii	Initial value for the EM algorithm. Each of them must be a vector of length g. (the algorithm considers the number of components to be adjusted based on the size of these vectors)
g	the number of cluster to be considered in fitting
get.init	if TRUE, the initial values are generated via k-means
criteria	It indicates if are calculated the criterion selection methods (AIC, BIC, EDC and ICL)

group	if TRUE, the vector with the classification of the response is returned
family	distribution family to be used in fitting ("Skew.t", "Skew.cn", "Skew.slash", "Skew.normal")
error	define the stopping criterion of the algorithm
iter.max	the maximum number of iterations of the EM algorithm
obs.prob	if TRUE, the posterior probability of each observation belonging to one of the g groups is reported
kmeans.param	a list with alternative parameters for the kmeans function when generating initial values, list(iter.max = 10, n.start = 1, algorithm = "Hartigan-Wong")
show.convergence	graphics of convergence for the parameters
cp	Cut Point

Value

The function returns a list with 16 elements detailed as

iter	Number of iterations.
criteria	Attained criteria value.
convergence	Convergence reached or not.
mu	Location parameter estimate.
sigma2	Scale parameter estimate.
lambda	Shape parameter estimate.
pii	Weight parameter estimate.
nu	Estimated degrees of freedom parameter.
SE	Standard Error estimates, if the output shows NA the function does not provide the standard error for this parameter.
table	Table containing the inference for the estimated parameters.
loglik	Log-likelihood value.
AIC	Akaike information criterion.
BIC	Bayesian information criterion.
EDC	Efficient Determination Criterion.
ICL	Information Completed Likelihood.
time	Processing time.

Author(s)

Luis Benites Sanchez <lbenitesanchez@gmail.com> and Rocio Paola Maehara <rmaeharaa@gmail.com> and Victor Hugo Lachos <hlachos@ime.unicamp.br>

References

- Basso, R. . M., Lachos, V. H., Cabral, C. R., Ghosh, P., 2010. Robust mixture modeling based on scale mixtures of skew-normal distributions. Computational Statistics & Data Analysis doi:10.1016/j.csda.2009.09.031.
- Lachos, V. H., Ghosh, P., Arellano-Valle, R. B., 2010. Likelihood based inference for skew - normal independent linear mixed models. Statistica Sinica 20, 303 - 322.

See Also

[FMsmnReg](#), [ais](#), [horses](#)

Examples

#See examples for the FMsmnReg function linked above.

horses

Horse Racing at Eagle Farm data

Description

Results of horse races at Eagle Farm, Brisbane, on 31 August 1998. The data, collected by Donald Forbes for his MS305 Data Analysis Project, give results for each horse in a sequence of 8 races.

Format

This data frame contains the following columns:

Position (Finishing position)

Starters Number of horses in race

Last Finishing position in last race

Since Days since last race

Number Identifying number of horse in race

Carried Weight carried

Weight Handicap weight

Barrier Barrier position at start of race

Distance Length of race

Lengths Number of lengths that horse finished from winner

Odds Starting odds

Starts Number of races previously started in

Age Age of horse in years

Ratio Proportion of wins in previous starts

References

Forbes, D. (1998). A Day at the Races. *MS305 Data Analysis Project, Department of Mathematics, University of Queensland.*

Examples

```
##Load the data
library(FMsmnReg)
data(horses)
attach(horses)

#Set the response y and covariate x
x1 <- cbind(1,Last,Carried)
y <- Position

##Fits a linear Regression Model with Finite Mixtures of Skew Contaminated Normal

parCN <- FMsmnReg(y, x1, g=2, get.init = TRUE, criteria = TRUE, group = FALSE,
  family = "Skew.cn", error = 10^-4, iter.max = 5000,obs.prob= FALSE,
  kmeans.param = NULL,show.converge=FALSE,cp=0.5)

##Fits a linear Regression Model with Finite Mixtures of Skew normal
parSN <- FMsmnReg(y, x1, g=2, get.init = TRUE, criteria = TRUE,
  group = FALSE,family = "Skew.normal", error = 10^-4,
  iter.max = 5000,obs.prob= FALSE, kmeans.param = NULL,show.converge=FALSE,cp=0.5)
```

Index

- * **EM**

- FMsmnReg, 4

- * **Finite Mixture**

- FMsmnReg, 4

- * **datasets**

- ais, 3

- horses, 6

- * **package**

- FMsmnReg, 4

- FMsmnReg-package, 2

- * **smsn**

- FMsmnReg, 4

ais, 3, 6

FMsmnReg, 2, 4, 6

FMsmnReg-package, 2

horses, 6, 6